

Gene Section

Mini Review

XRCC3 (X-ray repair complementing defective repair in Chinese hamster cells 3)

Ulla Vogel

National Research Centre for the Working Environment, Lerso Parkalle 105, DK-2100 Copenhagen O, Denmark

Published in Atlas Database: November 2005

Online updated version: <http://AtlasGeneticsOncology.org/Genes/XRCC3ID335ch14q32.html>
DOI: 10.4267/2042/38297

This work is licensed under a Creative Commons Attribution-Non-commercial-No Derivative Works 2.0 France Licence.
© 2006 Atlas of Genetics and Cytogenetics in Oncology and Haematology

Identity

Hugo: XRCC3
Location: 14q32.3

DNA/RNA

Description

17 800 bp, 9 exons.

Transcription

2,504 bp.

Protein

Description

346 amino acids.

Function

XRCC3 is required for efficient repair of double strand breaks via homologous recombinational repair (link), for correct chromosomal segregation and for repair of DNA cross links. Inactivation of XRCC3 in CHO cells results in increased radiation and cisplatin sensitivity. In human cells, XRCC3 forms a complex with Rad51C which is recruited early to DNA damage. Inactivation of XRCC3 in human cells leads to a two-fold sensitivity to DNA cross-linking agents, impaired Rad51 focus formation, elevated chromosome aberrations and five to seven-fold increased endoreduplication.

Homology

XRCC3 is a paralog to rad51.

Implicated in

No human disease has been linked to inactivation of XRCC3. However, polymorphisms in XRCC3 may be associated with increased cancer risk (see below).

Note: Genetic Epidemiology:

The most frequent polymorphism in XRCC3 is XRCC3 C18067T which results in a Thr to Met amino acid substitution at codon 241. Carriers of the variant T-allele of XRCC3 T241M have higher DNA adduct levels in lymphocyte DNA compared to homozygous C-allele carriers, indicating that the polymorphism is associated with lowered DNA repair capacity. The variant allele of XRCC3 T241M polymorphism has been associated with increased risk of squamous cell carcinoma of the head and neck in one study while another study found no association. No association has previously been found with colon cancer, non-melanoma skin cancer, prostate cancer, gastric cancer, ovarian cancer. Conflicting results have been published on the association with breast cancer, bladder cancer, malignant melanoma and lung cancer.

Two frequent SNP are upstream of XRCC3 T241M have also been studied, namely XRCC3 A4541G and XRCC3 A17897G. Neither polymorphism gives rise to amino acid changes. The variant allele of A17897G has been associated with decreased risk of breast cancer and ovarian cancer. The same tendency, but no significant associations was found for lung cancer. XRCC3 A4541G was not associated to risk of breast cancer or lung cancer. However, homozygote carriers of the variant allele had lower risk of serous epithelial ovarian cancer.

References

- Liu N, Lamerdin JE, Tebbs RS, Schild D, Tucker JD, Shen MR, Brookman KW, Siciliano MJ, Walter CA, Fan W, Narayana LS, Zhou ZQ, Adamson AW, Sorensen KJ, Chen DJ, Jones NJ, Thompson LH. XRCC2 and XRCC3, new human Rad51-family members, promote chromosome stability and protect against DNA cross-links and other damages. *Mol Cell* 1998;1(6):783-793.
- Griffin CS, Simpson PJ, Wilson CR, Thacker J. Mammalian recombination-repair genes XRCC2 and XRCC3 promote correct chromosome segregation. *Nat Cell Biol* 2000;2(10):757-761.
- Winsey SL, Haldar NA, Marsh HP, Bunce M, Marshall SE, Harris AL, Wojnarowska F, Welsh KI. A variant within the DNA repair gene XRCC3 is associated with the development of melanoma skin cancer. *Cancer Res* 2000;60(20):5612-5616.
- David-Beabes GL, Lunn RM, London SJ. No Association between the XPD (Lys751Gln) Polymorphism or the XRCC3 (Thr241Met) Polymorphism and Lung Cancer Risk. *Cancer Epidemiol Biomarkers Prev* 2001;10(8):911-912.
- Matullo G, Guarrera S, Carturan S, Peluso M, Malaveille C, Davico L, Piazza A, Vineis P. DNA repair gene polymorphisms, bulky DNA adducts in white blood cells and bladder cancer in a case-control study. *Int J Cancer* 2001a;92(4):562-567.
- Matullo G, Palli D, Peluso M, Guarrera S, Carturan S, Celentano E, Krogh V, Munnia A, Tumino R, Polidoro S, Piazza A, Vineis P. XRCC1, XRCC3, XPD gene polymorphisms, smoking and (32)P-DNA adducts in a sample of healthy subjects. *Carcinogenesis* 2001b;22(9):1437-1445.
- Duan Z, Shen H, Lee JE, Gershenwald JE, Ross MI, Mansfield PF, Duvic M, Strom SS, Spitz MR, Wei Q. DNA repair gene XRCC3 241Met variant is not associated with risk of cutaneous malignant melanoma. *Cancer Epidemiol Biomarkers Prev* 2002;11(10 Pt 1):1142-1143.
- Kuschel B, Auranen A, McBride S, Novik KL, Antoniou A, Lipscombe JM, Day NE, Easton DF, Ponder BA, Pharoah PD, Dunning A. Variants in DNA double-strand break repair genes and breast cancer susceptibility. *Hum Mol Genet* 2002;11(12):1399-1407.
- Shen H, Sturgis EM, Dahlstrom KR, Zheng Y, Spitz MR, Wei Q. A variant of the DNA repair gene XRCC3 and risk of squamous cell carcinoma of the head and neck: a case-control analysis. *Int J Cancer* 2002;99(6):869-872.
- Stern MC, Umbach DM, Lunn RM, Taylor JA. DNA repair gene XRCC3 codon 241 polymorphism, its interaction with smoking and XRCC1 polymorphisms, and bladder cancer risk. *Cancer Epidemiol Biomarkers Prev* 2002;11(9):939-943.
- Han J, Hankinson SE, Ranu H, De V, I, Hunter DJ. Polymorphisms in DNA double strand break repair genes and breast cancer risk in the nurses' health study. *Carcinogenesis* 2004;25(2):189-195.
- Jacobsen NR, Nexø BA, Olsen A, Overvad K, Wallin H, Tjønneland A, Vogel U. No association between the DNA repair gene XRCC3 T241M polymorphism and risk of skin cancer and breast cancer. *Cancer Epidemiol Biomarkers Prev* 2003;12(6):584-585.
- Mort R, Mo L, McEwan C, Melton DW. Lack of involvement of nucleotide excision repair gene polymorphisms in colorectal cancer. *Br J Cancer* 2003;89(2):333-337.
- Sanyal S, Festa F, Sakano S, Zhang Z, Steineck G, Norming U, Wijkström H, Larsson P, Kumar R, Hemminki K. Polymorphisms in DNA repair and metabolic genes in bladder cancer. *Carcinogenesis* 2004;25(5):729-734.
- Shen M, Hung RJ, Brennan P, Malaveille C, Donato F, Placidi D, Carta A, Hautefeuille A, Boffetta P, Porru S. Polymorphisms of the DNA repair genes XRCC1, XRCC3, XPD, interaction with environmental exposures, and bladder cancer risk in a case-control study in northern Italy. *Cancer Epidemiol Biomarkers Prev* 2003;12(11 Pt 1):1234-1240.
- Smith TR, Miller MS, Lohman K, Lange EM, Case LD, Mohrenweiser HW, Hu JJ. Polymorphisms of XRCC1 and XRCC3 genes and susceptibility to breast cancer. *Cancer Lett* 2003;190(2):183-190.
- Forget AL, Bennett BT, Knight KL. Xrcc3 is recruited to DNA double strand breaks early and independent of Rad51. *J Cell Biochem* 2004;93(3):429-436.
- Jacobsen NR, Raaschou-Nielsen O, Nexø B, Wallin H, Overvad K, Tjønneland A, Vogel U. XRCC3 polymorphisms and risk of lung cancer. *Cancer Lett* 2004;213(1):67-72.
- Popanda O, Schattenberg T, Phong CT, Butkiewicz D, Risch A, Edler L, Kayser K, Dienemann H, Schulz V, Drings P, Bartsch H, Schmezer P. Specific combinations of DNA repair gene variants and increased risk for non-small cell lung cancer. *Carcinogenesis* 2004;25(12):2433-2441.
- Yoshihara T, Ishida M, Kinomura A, Katsura M, Tsuruga T, Tashiro S, Asahara T, Miyagawa K. XRCC3 deficiency results in a defect in recombination and increased endoreduplication in human cells. *EMBO J* 2004;23(3):670-680.
- Auranen A, Song H, Waterfall C, Dicioccio RA, Kuschel B, Kjaer SK, Hogdall E, Hogdall C, Stratton J, Whittemore AS, Easton DF, Ponder BA, Novik KL, Dunning AM, Gayther S, Pharoah PD. Polymorphisms in DNA repair genes and epithelial ovarian cancer risk. *Int J Cancer* 2005;117(4):611-618.
- Festa F, Kumar R, Sanyal S, Undén B, Nordfors L, Lindholm B, Snellman E, Schalling M, Försti A, Hemminki K. Basal cell carcinoma and variants in genes coding for immune response, DNA repair, folate and iron metabolism. *Mutat Res* 2005;574(1-2):105-111.
- Huang WY, Chow WH, Rothman N, Lissowska J, Llaça V, Yeager M, Zatonski W, Hayes RB. Selected DNA repair polymorphisms and gastric cancer in Poland. *Carcinogenesis* 2005a;26(8):1354-1359.
- Huang WY, Olshan AF, Schwartz SM, Berndt SI, Chen C, Llaça V, Chanock SJ, Fraumeni JF, Jr., Hayes RB. Selected genetic polymorphisms in MGMT, XRCC1, XPD, and XRCC3 and risk of head and neck cancer: a pooled analysis. *Cancer Epidemiol Biomarkers Prev* 2005b;14(7):1747-1753.
- Raaphorst GP, Leblanc M, Li LF. A comparison of response to cisplatin, radiation and combined treatment for cells deficient in recombination repair pathways. *Anticancer Res* 2005;25(1A):53-58.
- Ritchey JD, Huang WY, Chokkalingam AP, Gao YT, Deng J, Levine P, Stanczyk FZ, Hsing AW. Genetic variants of DNA repair genes and prostate cancer: a population-based study. *Cancer Epidemiol Biomarkers Prev* 2005;14(7):1703-1709.
- Webb PM, Hopper JL, Newman B, Chen X, Kelemen L, Giles GG, Southey MC, Chenevix-Trench G, Spurdle AB. Double-strand break repair gene polymorphisms and risk of breast or ovarian cancer. *Cancer Epidemiol Biomarkers Prev* 2005;14(2):319-323.
- Yeh CC, Sung FC, Tang R, Chang-Chieh CR, Hsieh LL. Polymorphisms of the XRCC1, XRCC3, & XPD genes, and colorectal cancer risk: a case-control study in Taiwan. *BMC Cancer* 2005;5(1):12.

This article should be referenced as such:

Vogel U. XRCC3 (X-ray repair complementing defective repair in Chinese hamster cells 3). *Atlas Genet Cytogenet Oncol Haematol*.2006;10(2):105-106.
